

NULL SETS OF HARMONIC MEASURE ON NTA DOMAINS: LIPSCHITZ APPROXIMATION REVISITED

MATTHEW BADGER

ABSTRACT. We show the David-Jerison construction of big pieces of Lipschitz graphs inside a corkscrew domain does not require its surface measure be upper Ahlfors regular. Thus we can study absolute continuity of harmonic measure and surface measure on NTA domains of locally finite perimeter using Lipschitz approximations. A partial analogue of the F. and M. Riesz Theorem for simply connected planar domains is obtained for NTA domains in space. As a consequence every Wolff snowflake has infinite surface measure.

1. INTRODUCTION

What are the minimal assumptions on the boundary of a domain $\Omega \subset \mathbb{R}^n$ that guarantee its harmonic measure ω and surface measure $\sigma = \mathcal{H}^{n-1} \llcorner \partial\Omega$ have the same null sets? When $n = 2$, for example, one has the classic result of F. and M. Riesz [14]. In the plane, a topological condition ($\partial\Omega$ is a Jordan curve) and a mild measure-theoretic condition ($\partial\Omega$ has finite length) imply harmonic measure vanishes exactly on sets of zero length.

Theorem A (F. and M. Riesz 1916). *Let $\Omega \subset \mathbb{R}^2$ be a simply connected domain, bounded by a Jordan curve. If $\mathcal{H}^1(\partial\Omega) < \infty$, then*

$$(1.1) \quad \omega(E) = 0 \Leftrightarrow \mathcal{H}^1(E) = 0 \quad \text{for every Borel set } E \subset \partial\Omega.$$

When one strengthens the hypothesis $\mathcal{H}^1(\partial\Omega) < \infty$ in Theorem A, the relationship witnessed between ω and σ is stronger than absolute continuity [10]. A Jordan curve $\partial\Omega$ is called a *chord-arc curve* if $\partial\Omega$ is a quasicircle and there exists a constant $C > 0$ such that $\mathcal{H}^1(\Delta(Q, r)) \leq Cr$ for all $Q \in \partial\Omega$ and $0 < r < \text{diam } \Omega$, where $\Delta(Q, r) = \partial\Omega \cap B(Q, r)$.

Theorem B (Lavrentiev 1936). *Let $\Omega \subset \mathbb{R}^2$ be a simply connected domain, bounded by a chord-arc curve. Then (1.1) holds and $\omega \in A_\infty(\sigma)$: there exists constants $0 < \delta < 1$ and $0 < \varepsilon < 1$ such that for every $\Delta = \Delta(Q, r)$,*

$$(1.2) \quad \sigma(E) \leq \delta\sigma(\Delta) \Rightarrow \omega(E) \leq \varepsilon\omega(\Delta) \quad \text{for every Borel set } E \subset \Delta.$$

An amusing fact is that the one-sided inequality (1.2) implies (1.1). Actually $\omega \in A_\infty(\sigma)$ if and only if $\sigma \in A_\infty(\omega)$; see [2] also for several equivalent definitions of A -infinity weights. For further discussion on harmonic measure in the plane, the reader should consult [6].

The situation in higher dimensions is more delicate. Ziemer [17] found a topological sphere $\Omega \subset \mathbb{R}^3$ whose boundary is 2-rectifiable and $\mathcal{H}^2(\partial\Omega) < \infty$, but whose harmonic

Date: July 14, 2010.

2000 Mathematics Subject Classification. 28A75, 31A15.

Key words and phrases. Harmonic measure, absolute continuity, big pieces of Lipschitz graphs, corkscrew condition, NTA domain, Hausdorff dimension, Wolff snowflake.

The author was partially supported by NSF grants DMS-0838212 and DMS-0856687.

measure is supported on a subset of zero area. This means that any analogue of Theorem A in space must impose extra non-topological conditions on $\partial\Omega$. In this paper, we show the class of NTA domains (recalled in §4) satisfy the forward direction of (1.1).

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n$ be NTA. If $\mathcal{H}^{n-1} \llcorner \partial\Omega$ is Radon (e.g. if $\mathcal{H}^{n-1}(\partial\Omega) < \infty$), then $\partial\Omega$ is $(n-1)$ -rectifiable and*

$$(1.3) \quad \omega(E) = 0 \Rightarrow \mathcal{H}^{n-1}(E) = 0 \quad \text{for every Borel set } E \subset \partial\Omega.$$

The proof of Theorem 1.1 that we present is based on the extension of Theorem B to $n \geq 3$ given by David and Jerison [4]. Let $\Omega \subset \mathbb{R}^n$ be a NTA domain and assume its surface measure is *Ahlfors regular*; that is, there exists a constant $C > 0$ such that

$$(1.4) \quad C^{-1}r^{n-1} \leq \mathcal{H}^{n-1}(\Delta(Q, r)) \leq Cr^{n-1} \quad \text{for all } Q \in \partial\Omega \text{ and } 0 < r < r_0.$$

Using the existence of $(n-1)$ -disks inside $B(Q, r) \cap \Omega$ and $B(Q, r) \setminus \Omega$ of radius $\geq c_0 r$ (a weaker property than the corkscrew conditions enjoyed by NTA domains) and (1.4), David and Jerison gave a geometric construction of Lipschitz domains $\Omega_L \subset B(Q, r) \cap \Omega$ such that $\mathcal{H}^{n-1}(\partial\Omega_L \cap \partial\Omega) \geq c_1 r^{n-1}$. In other words, there exist Lipschitz approximations of Ω at each location and scale, with substantial intersection in the boundary. Applying Dahlberg's theorem relating harmonic and surface measures on Lipschitz domains [3] and a localization principle for NTA domains [8] yields Theorem B for NTA domains. (Theorem C was independently verified for 2-sided NTA domains by Semmes [15] using a stopping time argument.)

Theorem C (David and Jerison, Semmes 1990). *Let $\Omega \subset \mathbb{R}^n$ be NTA. If (1.4) holds, then $\omega \ll \sigma \ll \omega$ and $\omega \in A_\infty(\sigma)$.*

The existence of big pieces of Lipschitz graphs implies that every NTA domain satisfying (1.4) is uniformly rectifiable; this notion of quantitative rectifiability is developed in [5]. For a class of domains with non-doubling harmonic measure, on which a variant of the A_∞ condition in Theorem C still holds, see Bennewitz and Lewis [1]. In the present work, we revisit David and Jerison's construction of Lipschitz approximations to Ω in the case that Ω is a corkscrew domain (e.g. when Ω is NTA). We make two key observations. First the surface measure of any corkscrew domain is automatically lower Ahlfors regular (Lemma 2.3). Second the construction of an approximation at a given location and scale does not require surface measure be upper Ahlfors regular. Therefore one may relax the assumption that (1.4) holds uniformly at all scales in Theorem C. This is our main result.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^n$ be NTA. Then the set*

$$(1.5) \quad A = \left\{ Q \in \partial\Omega : \liminf_{r \downarrow 0} \frac{\mathcal{H}^{n-1}(\Delta(Q, r))}{r^{n-1}} < \infty \right\}$$

is $(n-1)$ -rectifiable and $\omega \llcorner A \ll \sigma \llcorner A \ll \omega \llcorner A$.

If $\mathcal{H}^{n-1} \llcorner \partial\Omega$ is Radon, then $\mathcal{H}^{n-1}(\partial\Omega \setminus A) = 0$. Hence Theorem 1.1 follows directly from Theorem 1.2. It is unknown whether the F. and M. Riesz theorem has a full analogue for NTA domains in higher dimensions. However, in view of Theorem 1.2, one can reverse the arrow in (1.3) if and only if

Conjecture 1.3. *Let $\Omega \subset \mathbb{R}^n$ be NTA. If $\mathcal{H}^{n-1} \llcorner \partial\Omega$ is Radon, then*

$$(1.6) \quad B = \left\{ Q \in \partial\Omega : \lim_{r \downarrow 0} \frac{\mathcal{H}^{n-1}(\Delta(Q, r))}{r^{n-1}} = \infty \right\}$$

has harmonic measure zero.

The paper is organized as follows. In sections 2–3, we show how to build a Lipschitz domain Ω_L inside of a corkscrew domain Ω at a given location $Q \in \partial\Omega$ and scale $r > 0$ such that $\mathcal{H}^{n-1}(\Delta(Q, r)) < \infty$. At each step of the construction, we keep careful track of dependencies on parameters. The common boundary $\partial\Omega \cap \partial\Omega_L$ of a domain and its approximation has size determined by the dimension n and corkscrew constant M of Ω ; the Lipschitz constant and character of an approximation only depends on n , M and the ratio $\gamma = \mathcal{H}^{n-1}(\Delta(Q, r))/r^{n-1}$. Section 2.1 outlines the construction of Ω_L using cones of a fixed aperture and reduces the approximation theorem (Theorem 2.4) to choosing the correct slope of the defining cones (Proposition 2.8). In section 2.2 we quantify the relationship between harmonic measure and surface measure on the Lipschitz domain Ω_L . The main tool is Jerison and Kenig’s proof [7] of Dahlberg’s theorem for star-shaped Lipschitz domains. The construction of Ω_L is completed in section 3, where we verify Proposition 2.8 by following the proof of the geometric proposition in [4].

Section 4 is devoted to the absolute continuity of harmonic measure on NTA domains. We derive Theorem 1.2 from three ingredients: the existence of good Lipschitz approximations to corkscrew domains (Theorem 2.4 and Lemma 2.13), a localization property of harmonic measure on NTA domains (Lemma 4.3), and a Vitali type covering theorem for Radon measures in \mathbb{R}^n (Theorem 4.6). An NTA domain is a corkscrew domain that satisfies a Harnack chain condition. The proof of absolute continuity that we give actually shows Theorem 1.2 is valid on any corkscrew domain whose harmonic measure satisfies the conclusion of Lemma 4.3.

Two applications of the main theorem are presented in section 5. First we demonstrate that every Wolff snowflake (studied in [16] and [11]) has infinite surface measure. Second we compute the Hausdorff dimension of harmonic measure on NTA domains of locally finite perimeter: if $\Omega \subset \mathbb{R}^n$ is NTA and $\mathcal{H}^{n-1} \llcorner \partial\Omega$ is Radon, then $\mathcal{H}\text{-dim } \omega = n - 1$. This section can be read independently of §§2–4.

2. LIPSCHITZ APPROXIMATION TO CORKSCREW DOMAINS

A closed set $\Sigma \subset \mathbb{R}^n$ has *big pieces of Lipschitz graphs* (abbreviated *BPLG*) if (i) $\mathcal{H}^{n-1} \llcorner \Sigma$ is Ahlfors regular and (ii) there are constants $\varphi > 0$, $h > 0$ and $r_0 > 0$ such that for every $Q \in \Sigma$ and $0 < r < r_0$ there exists (up to an isometry of \mathbb{R}^n) a graph $\Gamma \subset \mathbb{R}^n$ of an h -Lipschitz function such that $\mathcal{H}^{n-1}(B(Q, r) \cap \Sigma \cap \Gamma) \geq \varphi r^{n-1}$. In [4], David and Jerison proved if $\Sigma \subset \mathbb{R}^n$ has Ahlfors regular surface measure and the open set $\mathbb{R}^n \setminus \Sigma$ satisfies a “two disk” condition, then Σ has BPLG. Reading their proof carefully reveals that the upper bound in the Ahlfors regularity condition (1.4) is not used to build the Lipschitz graphs Γ . We verify this claim over the next two sections, in the special case that $\Sigma = \partial\Omega$ and $\Omega \subset \mathbb{R}^n$ is a corkscrew domain. (This is the case applicable for Theorem 1.2 and using the corkscrew condition instead of the two disk condition shortens the proof of several lemmas in §3.)

Definition 2.1. An open set $\Omega \subset \mathbb{R}^n$ satisfies the *corkscrew condition* with constants $M > 1$ and $R > 0$ provided for every $Q \in \partial\Omega$ and $0 < r < R$ there exists a *non-tangential point* $A = A(Q, r) \in \Omega$ such that $M^{-1}r < |A - Q| < r$ and $\text{dist}(A, \partial\Omega) > M^{-1}r$.

Definition 2.2. An open set $\Omega \subset \mathbb{R}^n$ is called a *corkscrew domain* if Ω is connected and both Ω and $\mathbb{R}^n \setminus \overline{\Omega}$ satisfy the corkscrew condition for some constants $M > 1$ and $R > 0$.

When $\Omega \subset \mathbb{R}^n$ is a corkscrew domain, we write $A^+(Q, r)$ for non-tangential points in the interior $\Omega^+ = \Omega$ and write $A^-(Q, r)$ for non-tangential points in the exterior $\Omega^- = \mathbb{R}^n \setminus \overline{\Omega}$. We do not require that the exterior of a corkscrew domain be connected.

Let us start with a simple application of the interior and exterior corkscrew conditions. Surface measure on a corkscrew domain is lower Ahlfors regular. Here and in the sequel, we normalize Hausdorff measure so that $\mathcal{H}^{n-1}(B_{n-1}(0, 1)) = \omega_{n-1}$.

Lemma 2.3. *There exists a constant $\beta = \beta(n, M) > 0$ such that for every corkscrew domain $\Omega \subset \mathbb{R}^n$ with constants $M > 1$ and $R > 0$,*

$$(2.1) \quad \mathcal{H}^{n-1}(\Delta(Q, r)) \geq \beta r^{n-1} \quad \text{for all } Q \in \partial\Omega \text{ and } r < R.$$

Proof. Write $t = M/(M + 1)$ and choose non-tangential points $a^\pm = A^\pm(Q, tr)$ of Ω^\pm . Then $B(a^\pm, tr/M) \subset \Omega^\pm \cap B(Q, r)$. Let π denote orthogonal projection onto a plane P (of codimension 1) orthogonal to the line segment connecting a^+ and a^- . Assign D^\pm to be the $(n - 1)$ -dimensional disk of radius tr/M inside of $B(a^\pm, tr/M)$ and parallel to P . Because D^+ and D^- lie in different components of $\mathbb{R}^n \setminus \partial\Omega$ and the ball $B(Q, r)$ is convex, any line segment from D^+ to D^- must intersect $\Delta(Q, r)$. Since $\pi(D^+) = \pi(D^-)$ is a disk of radius $tr/M = r/(M + 1)$,

$$(2.2) \quad \mathcal{H}^{n-1}(\Delta(Q, r)) \geq \mathcal{H}^{n-1}(\pi(\Delta(Q, r))) \geq \mathcal{H}^{n-1}(\pi(D^\pm)) = \omega_{n-1} \left(\frac{r}{M + 1} \right)^{n-1}.$$

Thus $\beta = \omega_{n-1}/(M + 1)^{n-1}$ suffices. □




FIGURE 1. Proof of Lemma 2.3

Our main goal in this section is to construct Lipschitz domains inside corkscrew domains with substantial intersection on the boundary. An important observation is the size of the big pieces of Lipschitz graphs depends only on the corkscrew constant.

Theorem 2.4. *Let $\Omega \subset \mathbb{R}^n$ be a corkscrew domain with constants $M > 1$ and $R > 0$. There exists a constant $\psi = \psi(n, M) > 0$ with the following property. For every $Q \in \partial\Omega$ and every positive number $r < R$ such that $\mathcal{H}^{n-1}(\Delta(Q, r)) < \infty$,*

- (\star) *for every non-tangential point $a = A^+(Q, r/2)$ there exists a Lipschitz domain $\Omega_L \subset \mathbb{R}^n$ such that $a \in \Omega_L \subset \Omega \cap B(Q, r)$, $\mathcal{H}^{n-1}(\partial\Omega_L \cap \partial\Omega) \geq \psi r^{n-1}$ and $\partial\Omega_L \cap \partial\Omega$ is contained in (the rigid motion of) a single Lipschitz graph.*

The Lipschitz constant and character of Ω_L depend only on n , M and $\mathcal{H}^{n-1}(\Delta(Q, r))/r^{n-1}$.

Corollary 2.5. *If $\Omega \subset \mathbb{R}^n$ is a corkscrew domain and $\mathcal{H}^{n-1} \llcorner \partial\Omega$ is upper Ahlfors regular, then $\partial\Omega$ has big pieces of Lipschitz graphs.*

Corollary 2.6. *Let $\Omega \subset \mathbb{R}^n$ be a corkscrew domain with constants $M > 1$ and $R > 0$. There exists a constant $h = h(n, M) > 0$ such that*

$$(2.3) \quad \{Q \in \partial\Omega : \mathcal{H}^{n-1}(\Delta(Q, r)) < \infty \text{ for some } r > 0\}$$

is contained, modulo a set of \mathcal{H}^{n-1} -measure zero, in the countable union of sets $F_i(\mathbb{R}^{n-1})$ where each function $F_i : \mathbb{R}^{n-1} \rightarrow \mathbb{R}^n$ has Lipschitz constant at most h .

2.1. Constructing a Lipschitz Approximation. Let $\Omega \subset \mathbb{R}^n$ be a corkscrew domain with constants $M > 1$ and $R > 0$, and fix $Q \in \partial\Omega$ and $0 < r < R$ such that

$$(2.4) \quad \mathcal{H}^{n-1}(\Delta(Q, r)) \leq \gamma r^{n-1} < \infty.$$

We do not normalize the radius $r = 1$, because we want to emphasize that the construction takes place at any fixed scale such that $\mathcal{H}^{n-1}(\Delta(Q, r))/r^{n-1} < \infty$. Our immediate goal is to find a constant $\psi = \psi(n, M) > 0$ such that (\star) in Theorem 2.4 holds.

Let a non-tangential point $a = A^+(Q, r/2)$ of Ω be given. We select a piece of the boundary to approximate as follows. Pick any non-tangential point $b = A^-(Q, r/2)$ of Ω^- . Then the line segment from a to b intersects $\Delta(Q, r/2)$ in some point Q' . After a harmless translation and rotation, we may assume that $Q' = 0$, $a = (0, a_n)$ and $b = (0, -b_n)$ where $r/2M \leq a_n, b_n \leq r$. Note that $B(a, r/2M) \subset B(Q, r) \cap \Omega$ and $B(b, r/2M) \subset B(Q, r) \cap \Omega^-$. Let $I_0 = [-s/2, s/2]^{n-1} \subset \mathbb{R}^{n-1}$ be the $(n-1)$ -dimensional cube with side length

$$(2.5) \quad s = \frac{r}{2M\sqrt{n-1}}$$

centered at the origin. Then $I_1 = I_0 \times \{a_n\} \subset B(a, r/4M)$. We shall write $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ for the orthogonal projections onto the first $(n-1)$ coordinates and the last coordinate of \mathbb{R}^n , respectively. Fix a cone \mathcal{C} opening upwards,

$$(2.6) \quad \mathcal{C} = \{z \in \mathbb{R}^n : f(z) \geq h|\pi(z)|\},$$

with parameter $h \gg 1$ to be chosen later. Let \mathcal{T} denote the trapezoidal region

$$(2.7) \quad \mathcal{T} = \{y \in \mathbb{R}^n : -b_n \leq f(y) \leq a_n \text{ and } (y + \mathcal{C}) \cap f^{-1}(a_n) \subset I_1\}$$

and write $T = \partial\Omega \cap \mathcal{T}$ for the portion of the boundary in \mathcal{T} . We shall approximate T . Since $I_0 \times [a_n - s/4, a_n] \subset B(a, r/2M) \subset \Omega$ and $I_0 \times [-b_n, -b_n + s/4] \subset B(b, r/2M) \subset \Omega^-$, we know that

$$(2.8) \quad -b_n + s/4 \leq f(y) \leq a_n - s/4 \quad \text{for all } y \in T.$$

Lemma 2.7. *If $h \geq h_0 = 16M\sqrt{n-1}$, then $\mathcal{H}^{n-1}(\pi(T)) \geq r^{n-1} / (4M\sqrt{n-1})^{n-1}$.*

Proof. If h is sufficiently large, then \mathcal{T} contains $I_2 = [-s/4, s/4]^{n-1} \times \{-b_n\}$. To check this claim, first note $I_2 \subset \mathcal{T}$ if and only if the corner $y = (s/4, \dots, s/4, -b_n)$ of the cube satisfies $(y + \mathcal{C}) \cap f^{-1}(a_n) \subset I_1$. Hence if $z \in \mathbb{R}^n$, $f(z) = h|\pi(z)|$ and $-b_n + f(z) = f(y + z) = a_n$ (i.e. $y + z \in (y + \partial\mathcal{C}) \cap f^{-1}(a_n)$) then $s/4 + |\pi(z)| \leq s/2$ (i.e. $y + z \in I_1$) implies $I_2 \subset \mathcal{T}$. That is,

$$(2.9) \quad \frac{s}{4} + \frac{a_n + b_n}{h} \leq \frac{s}{2} \quad \Rightarrow \quad I_2 \subset \mathcal{T}.$$

Since $a_n + b_n \leq 2r = 4Ms\sqrt{n-1}$, we find that $I_2 \subset \mathcal{T}$ provided

$$(2.10) \quad \frac{s}{4} + \frac{4Ms\sqrt{n-1}}{h} \leq \frac{s}{2}.$$

Thus $I_2 \subset \mathcal{T}$ when $h \geq h_0 = 16M\sqrt{n-1}$. Because every vertical line segment from $I_2 \subset \Omega^- \cap \mathcal{T}$ to $I_1 \subset \Omega^+ \cap \mathcal{T}$ intersects T and $\pi(I_2) \cap \pi(I_1) = \pi(I_2)$, we conclude

$$(2.11) \quad \mathcal{H}^{n-1}(\pi(T)) \geq \mathcal{H}^{n-1}(\pi(I_2)) = \left(\frac{s}{2}\right)^{n-1} = \left(\frac{r}{4M\sqrt{n-1}}\right)^{n-1}$$

whenever $h \geq h_0$. □




FIGURE 2. What do T_Γ and Ω_L look like?

We now use the cone \mathcal{C} to identify a subset of T which intersects a Lipschitz graph inside the closure of $\Omega \cap B(Q, r)$ in a big piece. By a standard argument the set T_Γ ,

$$(2.12) \quad T_\Gamma = \{y \in T : (y + \mathcal{C}) \cap T = \{y\}\},$$

is contained in the graph Γ of a function $F : I_0 \rightarrow \mathbb{R}$ with Lipschitz constant at most h . By (2.8) replacing F by $\max(\min(F, a_n - s/4), -b_n + s/4)$ does not effect $\Gamma \cap T$. Hence we may assume without loss of generality that $-b_n + s/4 \leq F(x) \leq a_n - s/4$ for all $x \in I_0$. Define the domain

$$(2.13) \quad \Omega_L = \{(x, u) \in I_0 \times \mathbb{R} : u > F(x)\} \cap \text{int}(\mathcal{T} \cup (I_0 \times [a_n, a_n + s/4])).$$

That is, Ω_L is obtained by taking the area above Γ inside \mathcal{T} and then extending upwards so that $B(a, s/4)$ lies inside the domain. Since the extension $I_0 \times [a_n, a_n + s/4] \subset B(a, r/2M)$, $\Omega_L \subset \Omega \cap B(0, r)$ and Ω_L is a Lipschitz domain with intersection $\partial\Omega_L \cap \partial\Omega = \Gamma \cap T = T_\Gamma$. Notice that Ω_L can be covered by $c(n)$ Lipschitz graphs with constant at most h .

To select the slope h of \mathcal{C} large enough so that $\mathcal{H}^{n-1}(T_\Gamma) \geq \psi r^{n-1}$ for some constant $\psi = \psi(n, M) > 0$, we need the following claim, a slight modification of the geometric proposition in [4].

Proposition 2.8. *For all $\varepsilon > 0$, there exists $h \geq h_0$ depending only on n , M , γ and ε such that $\mathcal{H}^{n-1}(\pi(T) \setminus \pi(T_\Gamma)) \leq \varepsilon r^{n-1}$.*

The proof of Proposition 2.8 is a long but fairly straightforward application of the corkscrew condition on the exterior Ω^- of Ω , the lower Ahlfors regularity of $\mathcal{H}^{n-1} \llcorner \partial\Omega$, and the upper bound $\mathcal{H}^{n-1}(\Delta(0, r)) \leq \gamma r^{n-1}$; details are postponed until §3. First let us finish studying the Lipschitz approximation Ω_L to $\Omega \cap B(Q, r)$.

Proof of Theorem 2.4. By Lemma 2.7 and Proposition 2.8,

$$(2.14) \quad \begin{aligned} \mathcal{H}^{n-1}(T_\Gamma) &\geq \mathcal{H}^{n-1}(\pi(T_\Gamma)) = \mathcal{H}^{n-1}(\pi(T)) - \mathcal{H}^{n-1}(\pi(T) \setminus \pi(T_\Gamma)) \\ &\geq \left(\frac{1}{(4M\sqrt{n-1})^{n-1}} - \varepsilon \right) r^{n-1}. \end{aligned}$$

Thus, choosing $\varepsilon > 0$ sufficiently small (equivalently choosing $h \geq h_0$ sufficiently large), we conclude that $\mathcal{H}^{n-1}(\partial\Omega_L \cap \partial\Omega) = \mathcal{H}^{n-1}(T_\Gamma) \geq \psi r^{n-1}$ with

$$(2.15) \quad \psi = \frac{1}{(8M\sqrt{n-1})^{n-1}}.$$

The constant ψ only depends on n and M ; the Lipschitz constant of the graph Γ and the Lipschitz character of the domain Ω_L are determined by h and thus by n , M and γ . \square

2.2. Harmonic Measure and the A -infinity Condition. Next we compare harmonic measure ω_L and surface measure σ_L on Ω_L (see Lemma 2.13) using constants depending only on n , M and γ . A theorem of Dahlberg [3] asserts a strong relationship between harmonic measure and surface measure exists on any bounded Lipschitz domain.

Theorem 2.9 ([3] Theorem 3). *Let $D \subset \mathbb{R}^n$ be a bounded Lipschitz domain equipped with harmonic measure ω_D and surface measure σ_D . Then $\omega_D \in A_\infty(\sigma_D)$.*

To use Theorem 2.9 effectively, we need to understand the dependence of the constants in the A_∞ condition on the features of a Lipschitz domain. There are currently two proofs of Theorem 2.9 (Dahlberg [3], Jerison and Kenig [7]) and each proof establishes the theorem first on a special class of star-shaped Lipschitz domains. (A domain D is called *star-shaped* with *center* c if every open line segment from c to ∂D lies inside D .)

Thus the A_∞ condition in Theorem 2.9 may depend on how star-shaped Lipschitz domains cover the original domain. To clarify this dependence, we need to introduce some notation.

Let D be a star-shaped Lipschitz domain with center c , and write \vec{n}_y for the outer unit normal to ∂D defined at \mathcal{H}^{n-1} -a.e. $y \in \partial D$. We define the *angle function* $\vartheta_{D,c}(y)$ by

$$(2.16) \quad \cos \vartheta = \frac{\langle \vec{n}_y, y - c \rangle}{\|y - c\|}, \quad 0 \leq \vartheta \leq \pi.$$

Note $\vartheta_{D,c}(y)$ is defined at \mathcal{H}^{n-1} -a.e. $y \in \partial D$. Also, since D is star-shaped with center c , the angle function $\vartheta_{D,c}(y) \in [0, \pi/2]$ almost surely.




FIGURE 3. The angle function $\vartheta_{D,c}(y)$

The following proposition is adapted from the proof of Theorem 2.9 in [7].

Proposition 2.10. *Let D be a bounded star-shaped Lipschitz domain with center $c \in D$. Let h be the Lipschitz constant of D and assume there exists radii $\rho_1, \rho_2 > 0$ such that $B(c, \rho_1) \subset D \subset B(c, \rho_2)$. For all $\vartheta_0 < \pi/2$, there exists a constant $C = C(n, h, \rho_2/\rho_1, \vartheta_0)$ with the following property. If $\vartheta_{D,c} \leq \vartheta_0$ a.e. on $\Delta_D(y_0, r_0)$ for some $y_0 \in \partial D$ and $r_0 > 0$, then the Radon-Nikodym derivative $k = d\omega_D^c/d\sigma_D$ of harmonic measure with pole at c with respect to surface measure satisfies the reverse Hölder inequality*

$$(2.17) \quad \left(\int_{\Delta(x,r)} k^2 d\sigma_D \right)^{1/2} \leq C \int_{\Delta(x,r)} k d\sigma_D \quad \text{for every } \Delta_D(y, r) \subset \Delta_D(y_0, r_0).$$

Here the dashed integral $\int_{\Delta} k d\sigma = (\sigma(\Delta))^{-1} \int_{\Delta} k d\sigma$ denotes an average. By the theory of A -infinity weights, if (2.17) holds, then (1.2) also holds on $\Delta_D(y_0, r_0)$ with constants δ and ε which only depend on n and C . Let us return our attention to the comparison of harmonic measure and surface measure on Ω_L .

First we cover Ω_L using two types of star-shaped Lipschitz domains (see Figure 4) and estimate harmonic measure in each case separately.

Lemma 2.11. *Up to a dilation, the Lipschitz domain*

$$(2.18) \quad D_{\text{top}} = \{y \in \Omega_L : f(y) > a_n - s/4\}$$

is determined by n and h . Moreover, there exists a constant $\eta_0 = \eta_0(n, M, \gamma)$ such that

$$(2.19) \quad \omega_L^a(E) \leq \eta_0 \Rightarrow \mathcal{H}^{n-1}(E) \leq \frac{\psi}{4} r^{n-1} \quad \text{for all } E \subset \partial\Omega_L \cap \partial D_{\text{top}}$$

Here ω_L^a denotes harmonic measure of Ω_L with pole at a .




FIGURE 4. Star-shaped domains in Lemmas 2.11 and 2.12

Proof. We required that the Lipschitz function $F : I_0 \rightarrow \mathbb{R}$ used to construct Ω_L satisfy $F(x) \leq a_n - s/4$ for all $x \in I_0$. Hence D_{top} is a fixed subset of the region $\text{int}(\mathcal{T} \cup (I_0 \times [a_n, a_n + s/4]))$ and determined by n and h (see Figure 3). By Theorem 2.9, we conclude $\omega_{\text{top}} \in A_\infty(\sigma_{\text{top}})$, where ω_{top}^a is harmonic measure on D_{top} with pole at a and $\sigma_{\text{top}} = \mathcal{H}^{n-1} \llcorner \partial D_{\text{top}}$. Thus there exist constants $p, q > 0$ depending only on n and h such that

$$(2.20) \quad \mathcal{H}^{n-1}(E) \leq p [\omega_{\text{top}}^a(E)]^q \mathcal{H}^{n-1}(D_{\text{top}}) \quad \text{for all } E \subset \partial D_{\text{top}}.$$

On one hand, $\omega_{\text{top}}^a(E) \leq \omega_L^a(E)$ for every $E \subset \partial \Omega_L \cap \partial D_{\text{top}}$ by the maximum principle. On the other hand, $\mathcal{H}^{n-1}(D_{\text{top}}) \leq C_0 r^{n-1}$ for some $C_0 = C_0(n, h)$. Therefore,

$$(2.21) \quad \mathcal{H}^{n-1}(E) \leq C_0 p [\omega_L^a(E)]^q r^{n-1} \quad \text{for all } E \subset \partial \Omega_L \cap \partial D_{\text{top}},$$

and the constant $\eta_0 = (\psi/4C_0p)^{1/q}$ depending only on n , M and γ suffices. \square

Lemma 2.12. Set $h^* = h\sqrt{n-1}$. For each $c \in I_1$ define

$$(2.22) \quad D_c = \{y \in \Omega_L : \|\pi(y) - \pi(c)\|_\infty < s/8h^*\}$$

where $\|x\|_\infty = \max_i |x_i|$ for all $x \in \mathbb{R}^{n-1}$. Assume $\|\pi(c)\|_\infty \leq s/2 - s/4h^* - s/8h^*$. Then D_c is a star-shaped Lipschitz domain with center c , the Lipschitz constant of D_c is at most h and $B(c, s/8h^*) \subset D_c \subset B(c, 4Ms\sqrt{n-1})$. Moreover, there exists a constant $\vartheta_1 = \vartheta_1(n, M, \gamma) < \pi/2$ such that $\vartheta_{D,c} \leq \vartheta_1$ on ∂D_c almost surely.

Proof. Fix any $c \in I_1$ such that $\|\pi(c)\|_\infty \leq s/2 - s/4h^* - s/8h^*$. Then $\|\pi(y)\|_\infty \leq s/2 - s/4h^*$ for all $y \in D_c$ and the top portion of D_c is a box that (up to translation) is independent of c :

$$(2.23) \quad D_c \cap D_{\text{top}} = (\pi(c) - s/8h^*, \pi(c) + s/8h^*)^{n-1} \times (a_n - s/4, a_n + s/4).$$

Hence $B(c, s/8h^*) \subset D_c$. The inclusion $D_c \subset B(c, 4Ms\sqrt{n-1})$ follows easily from (2.8). The bottom portion of D_c (i.e. $D_c \cap f^{-1}(-\infty, a_n - s/4)$) is the area above the graph

$$(2.24) \quad \Gamma_c = \{y \in \partial \Omega_L \cap \Gamma : \|\pi(y) - \pi(c)\|_\infty < s/8h^*\}.$$

For all $p > 0$ let $\mathcal{C}_p = \{z \in \mathbb{R}^n : f(z) \geq p|\pi(z)|\}$ denote the cone opening upwards with slope p . The cone used to construct Ω_L above was $\mathcal{C} = \mathcal{C}_h$. If $y \in \Gamma_c$, then $c \in y + \mathcal{C}_{2h}$ since $f(c) - f(y) = a_n - f(y) \geq s/4$ and $|\pi(c) - \pi(y)| \leq s/8h$. Because $\mathcal{C}_{2h} \subset \mathcal{C}_h$ and $(y + \mathcal{C}_h) \cap \Gamma = \{y\}$, the open segment from c to $y \in \Gamma_c$ lies in D_c . Thus D_c is star-shaped with respect to c . It remains to bound the angle function.

Suppose \vec{n}_y is an outer normal to ∂D_c defined at $y \in \Gamma_c$. Since $(y + \mathcal{C}_h) \cap \Gamma = \{y\}$, we know that $\vec{n}_y \in -\mathcal{C}_{1/h}$. On the other hand, we also know that $y - c \in -\mathcal{C}_{2h}$. The greatest angle between a vector $\vec{v} \in -\mathcal{C}_{1/h}$ and a vector $\vec{w} \in -\mathcal{C}_{2h}$ is obtained by $\vec{v} = (1, 0, \dots, -1/h)$ and $\vec{w} = (-1, 0, \dots, -2h)$; in this case,

$$(2.25) \quad \cos \vartheta = \frac{\langle \vec{v}, \vec{w} \rangle}{\|\vec{v}\| \|\vec{w}\|} = \frac{1}{(1 + h^{-2})^{1/2} (1 + 4h^2)^{1/2}} \geq \frac{1}{h\sqrt{10}}.$$

We conclude that $\vartheta_{D,c}(y) \leq \cos^{-1}(1/h\sqrt{10}) < \pi/2$ for almost every $y \in \Gamma_c$. Bounding $\vartheta_{D,c}$ on $\partial D_c \setminus \Gamma_c$ (that is, on sides of a box) is easy and left to the reader. \square

Equipped with Proposition 2.10, Lemma 2.11 and Lemma 2.12, we are ready to compare harmonic measure and surface measure on Ω_L .

Lemma 2.13. *There exists a constant $0 < \eta < 1$ depending only on n , M and γ with the following property. For every Borel set $E \subset \partial\Omega_L$,*

$$(2.26) \quad \omega_L^a(E) \leq \eta \Rightarrow \mathcal{H}^{n-1}(E) \leq \frac{\psi}{2} r^{n-1}.$$

Here ω_L^a denotes harmonic measure on Ω_L with pole at a .

Proof. Choose points $c_1, \dots, c_{i_0} \in I_1$ such that $\|\pi(c_i)\|_\infty \leq s/2 - s/4h^* - s/8h^*$ and

$$(2.27) \quad (-s/2 + s/4h^*, s/2 - s/4h^*)^{n-1} \times \{a_n\} \subset \bigcup_i D_{c_i}.$$

We can make this choice so that i_0 only depends on n and h . Notice that

$$(2.28) \quad \partial\Omega_L \subset \partial D_{\text{top}} \cup \bigcup_i \Gamma_{c_i}.$$

The points c_i and a lie inside D_{top} , at a uniform distance away from the boundary. Hence by Lemma 2.11 and Harnack's inequality there exists a constant $C_1 = C_1(n, h) > 1$ such that

$$(2.29) \quad \omega_L^{c_i}(E) \leq C_1 \omega^a(E) \quad \text{for all } 1 \leq i \leq i_0 \text{ and } E \subset \partial\Omega_L.$$

Thus, in view of (2.19), (2.28) and (2.29), to prove Lemma 2.13 it suffices to display $\eta = \eta(n, M, \gamma) \in (0, \eta_0)$ small enough so that

$$(2.30) \quad \omega_L^{c_i}(E \cap \Gamma_{c_i}) \leq C_1 \eta \Rightarrow \mathcal{H}^{n-1}(E \cap \Gamma_{c_i}) \leq \frac{\psi}{4i_0} r^{n-1}$$

for all $1 \leq i \leq i_0$ and $E \subset \partial\Omega$.

By Proposition 2.10 and Lemma 2.12, D_{c_i} is a star-shaped Lipschitz domain whose harmonic measure $\omega_D^{c_i}$ satisfies (2.17) on every disk for some constant C_2 depending only on n , M and γ . An equivalent form of the A -infinity condition is that for every $\varepsilon > 0$ there exists $\delta = \delta(n, C_2, \varepsilon) > 0$ such that

$$(2.31) \quad \omega_D^{c_i}(E \cap \Gamma_{c_i}) \leq \delta \Rightarrow \mathcal{H}^{n-1}(E \cap \Gamma_{c_i}) \leq \varepsilon \mathcal{H}^{n-1}(\partial D_{c_i}).$$

But $\mathcal{H}^{n-1}(\partial D_{c_i}) \leq C_3(n, h)r^{n-1}$, so choosing $\varepsilon = C_3^{-1}\psi/4i_0$ there exists $\delta = \delta(n, M, \gamma) > 0$ such that

$$(2.32) \quad \omega_D^{c_i}(E \cap \Gamma_{c_i}) \leq \delta \Rightarrow \mathcal{H}^{n-1}(E \cap \Gamma_{c_i}) \leq \frac{\psi}{4i_0}r^{n-1}.$$

Set $\eta = \min(\eta_0, \delta/C_1)$ so that η only depends on n, M and γ . Then (2.30) follows from (2.32) and the maximum principle. \square

3. PROOF OF PROPOSITION 2.8

We continue to assume the notation adopted in §2.1. Let $\varepsilon > 0$ be given. Our goal is to choose the slope $h \geq h_0$ of the cone \mathcal{C} so that $\mathcal{H}^{n-1}(\pi(T) \setminus \pi(T_\Gamma)) \leq \varepsilon r^{n-1}$. In the course of exposition we shall introduce several constants and indicate their dependence on previously defined quantities; each one will ultimately depend on at most n (dimension), M (corkscrew condition), γ (upper bound on scale r) and ε . To start we break up the set $\pi(T) \setminus \pi(T_\Gamma)$ into manageable pieces.

Lemma 3.1. *Let $H : \mathbb{R}^{n-1} \rightarrow [0, \infty]$ be the function*

$$(3.1) \quad H(x) = \sup \left\{ \frac{\mathcal{H}^{n-1}(\Delta(Q, r) \cap \pi^{-1}(I))}{\mathcal{H}^{n-1}(I)} : I \subset \mathbb{R}^{n-1} \text{ is a cube and } x \in I \right\}.$$

If $\lambda_H(N) = \{x \in \mathbb{R}^{n-1} : H(x) \geq N\}$, then $\mathcal{H}^{n-1}(\lambda_H(N)) \leq 5^{n-1}\gamma r^{n-1}/N$.

Proof. Just note $H(x)$ is the maximal function of the the measure

$$(3.2) \quad \mu = \pi_\#(\mathcal{H}^{n-1} \llcorner \Delta(Q, r))$$

with respect to Lebesgue measure on \mathbb{R}^{n-1} . By the Hardy-Littlewood maximal theorem (for example, see Theorem 2.19 in [13]), $\mathcal{H}^{n-1}(\lambda_H(N)) \leq 5^{n-1}\mu(\mathbb{R}^{n-1})/N$. By (2.4), μ has total mass $\mu(\mathbb{R}^{n-1}) \leq \gamma r^{n-1}$. \square

Note that the upper bound $\mathcal{H}^{n-1}(\Delta(Q, r)) \leq \gamma r^{n-1}$ on surface measure was used in the proof of Lemma 3.1. It will not be used again until after proof of Lemma 3.8.

For the remainder of the proof, fix $N = 2 \cdot 5^{n-1}\gamma/\varepsilon$ so that the set $\Lambda = \lambda_H(N)$ of points in \mathbb{R}^{n-1} where the maximal function $H(x)$ is large has small measure, $\mathcal{H}^{n-1}(\Lambda) \leq (\varepsilon/2)r^{n-1}$. We need to control the size of $\pi(T) \setminus \pi(T_\Gamma)$ in Λ^c . For each $y \in \mathcal{T}$, let $L(y) = [y, (\pi(y), a_n)]$ denote the vertical segment above y in \mathcal{T} . Then the set of points


$$(3.3) \quad T_E = \{y \in T : L(y) \cap T = \{y\}\}$$

denotes the “top edge” of $\partial\Omega$ inside \mathcal{T} . Observe that $T_\Gamma \subset T_E \subset T$ and $\pi(T) = \pi(T_E)$. Let α be a large power of 2 to be chosen later (after Lemma 3.8) and abbreviate $s_p = s/\alpha^p$ for all p . For each integer $k \geq 0$, define the set $F_k \subset I_0$,

$$(3.4) \quad F_k = \{\pi(y) : \exists y, z \in T_E \text{ such that } z \in y + \mathcal{C} \text{ and } s_k \leq f(z) - f(y) \leq s_{k-1}\}.$$

Since $-b_n + s/4 \leq f(y) \leq a_n - s/4$ for all $y \in T$, if we choose $\alpha s = s_{-1} \geq a_n + b_n - s/2$ then each bad point $x \in \pi(T) \setminus \pi(T_\Gamma)$ belongs to at least one F_k . For this reason, we will stipulate that

$$(3.5) \quad \alpha \geq 4M\sqrt{n-1} = 2r/s > s^{-1}(a_n + b_n - s/2).$$

FIGURE 5. A point $\pi(y) \in F_k$

Thus, we can prove the proposition by verifying $\mathcal{H}^{n-1}(\bigcup_k F_k) \leq \varepsilon r^{n-1}$, or by Lemma 3.1, by demonstrating that

$$(3.6) \quad \sum_k \mathcal{H}^{n-1}(F_k \setminus \Lambda) \leq \frac{\varepsilon}{2} r^{n-1} \quad \text{for some } h \geq h_0.$$

Remark 3.2. We do not assert that T_E or F_k are measurable. While this fact is irksome, it does not hinder the proof. A careful reader will observe that we only use countable subadditivity of the outer measure \mathcal{H}^{n-1} in coverings involving F_k .

The next lemma captures a simple idea. If the surface $\partial\Omega \cap \pi^{-1}(I)$ over a cube $I \subset \mathbb{R}^{n-1}$ has a big vertical span relative to the width of I , then the maximal function is big on I (by lower Ahlfors regularity). Let $\beta = \beta(n, M)$ be the constant given in Lemma 2.3. Also define the n -dimensional box $\mathcal{R} = 2I_0 \times [-b_n, a_n]$ and note $\mathcal{T} \subset \mathcal{R} \subset B(Q, r)$.

Lemma 3.3. *Let $I \subset 2I_0$ be a cube of side length t . Suppose one can find line segments L in $\Omega \cap \mathcal{R}$ and L' in $\Omega^- \cap \mathcal{R}$ such that $\pi(L)$ and $\pi(L')$ belong to I and $f(L) \cap f(L')$ is a segment of length $\geq \lceil 4^{n-1} \beta^{-1} N \rceil t$. Then the cube I belongs to Λ .*

Proof. Select $\lceil 4^{n-1} \beta^{-1} N \rceil$ points u_i in $f(L) \cap f(L')$ such that $|u_i - u_j| \geq t$ for $i \neq j$. Then for each u_i the horizontal line segment in $f^{-1}(u_i)$ which joins L to L' intersects $\partial\Omega$ at some point y_i , because L and L' belong to different components of $\mathcal{R} \setminus \partial\Omega$. The balls $\Delta(y_i, t/2)$ are disjoint sets and by Lemma 2.3 (note $t \ll 2s < r < R$),

$$(3.7) \quad \mathcal{H}^{n-1} \left(\bigcup_i \Delta(y_i, t/2) \right) \geq (4^{n-1} \beta^{-1} N) \beta \left(\frac{t}{2} \right)^{n-1} = N \mathcal{H}^{n-1}(2I).$$

Hence, $2I \subset \Lambda$, since $\Delta(y_i, t/2) \subset \Delta(0, r) \cap \pi^{-1}(2I)$ for each y_i (one easily checks that the distance of \mathcal{R} to $\mathbb{R}^n \setminus B(0, r)$ is much farther than $t/2$). In particular, $I \subset \Lambda$. \square

It will be convenient to work with a “dyadic” decomposition of I_0 . We say that a cube $I \subset I_0$ is *admissible* if the cube $\{(1/2, \dots, 1/2) + x/s : x \in I\}$ is dyadic in the usual sense. Hence I_0 is admissible and every admissible cube $I \subset I_0$ of side length $s/2^i$ is the almost

disjoint union of $2^{j(n-1)}$ admissible cubes of side length $s/2^{i+j}$. Moreover, given any cube $I \subset I_0$ of side length t , we can find an admissible cube contained in I of side length $\geq t/4$. The following “search lemma” locates admissible cubes inside $\Lambda \cup F_k^c$.

Lemma 3.4. *There exist constants $C_1 = C_1(n, N, \alpha, \beta)$ and $C_2 = C_2(n, M, N, \beta)$ with the following property. If $I \subset I_0$ is any cube with side length t satisfying*

$$(3.8) \quad \frac{C_1}{h} s_k \leq t \leq s_k,$$

then one can find an admissible cube $J \subset I \cap (\Lambda \cup F_k^c)$ of side length $\geq t/C_2$.

Proof. Let $I \subset I_0$ with side length t , $C_1 h^{-1} s_k \leq t \leq s_k$ be given. Since any cube contains an admissible cube of comparable size, we may first search for a cube $J \subset I \cap (\Lambda \cup F_k^c)$ which is not necessarily admissible.

For every cube $\mathcal{Q} \subset \mathbb{R}^{n-1}$ such that $\pi(T) \cap \mathcal{Q} \neq \emptyset$, define

$$(3.9) \quad \lambda(\mathcal{Q}) = \max\{f(y) : y \in T_E \text{ and } \pi(y) \in \mathcal{Q}\}.$$

The maximum in (3.9) is realized, because T_E is the “top edge” of T and T is compact. Suppose that there exists a cube $\mathcal{Q} \subset I$ of side length qt such that $\lambda(\mathcal{Q}) \leq \lambda(\frac{1}{2}\mathcal{Q}) + s_k/8$. Pick $w \in T_E$ such that $\pi(w) \in \frac{1}{2}\mathcal{Q}$ and $f(w) = \lambda(\frac{1}{2}\mathcal{Q})$, and let $c = A^-(w, qt/8)$ be any non-tangential point of Ω^- . Then $B(c, qt/8M) \subset \Omega^-$. We assign $K \subset \mathbb{R}^{n-1}$ to be the $(n-1)$ -dimensional cube with center $\pi(c)$ and side length t_K ,

$$(3.10) \quad t_K = \lceil 4^{n-1} \beta^{-1} N \rceil^{-1} qt/4M.$$

Note that $K \subset \mathcal{Q} \subset I$, since $\pi(c)$ is the center of K , $\pi(w) \in \frac{1}{2}\mathcal{Q}$, $\text{dist}(\frac{1}{2}\mathcal{Q}, \mathcal{Q}^c) = qt/4$ and

$$(3.11) \quad \text{dist}(x, \frac{1}{2}\mathcal{Q}) \leq \frac{\text{diam } K}{2} + |\pi(c) - \pi(w)| \ll \frac{qt}{16} + \frac{qt}{8} = \frac{3qt}{16} \quad \text{for all } x \in K.$$

If $K \cap F_k = \emptyset$, we are done. Otherwise there exist points $y, z \in T_E$ such that $\pi(y) \in K$, $z \in y + \mathcal{C}$ and $s_k \leq f(z) - f(y) \leq s_{k-1}$. We claim that $\pi(z) \in \mathcal{Q}$ if C_1 is sufficiently large. Because $z \in y + \mathcal{C}$ and $C_1 h^{-1} s_k \leq t$,

$$(3.12) \quad |\pi(z) - \pi(y)| \leq h^{-1}(f(z) - f(y)) \leq h^{-1} \alpha s_k \leq t \alpha / C_1.$$

Since $\pi(y) \in K$, using (3.11) it follows that

$$(3.13) \quad \text{dist}(\pi(z), \frac{1}{2}\mathcal{Q}) \leq |\pi(z) - \pi(y)| + \text{dist}(\pi(y), \frac{1}{2}\mathcal{Q}) \leq t \left(\frac{\alpha}{C_1} + \frac{3q}{16} \right).$$

Hence, $\text{dist}(\pi(z), \frac{1}{2}\mathcal{Q}) \leq qt/4$ and $\pi(z) \in \mathcal{Q}$ provided that $C_1 \geq 16\alpha/q$. Assume that C_1 has been chosen so that this is true. Then

$$(3.14) \quad f(y) \leq f(z) - s_k \leq \lambda(\mathcal{Q}) - s_k \leq \lambda(\frac{1}{2}\mathcal{Q}) - \frac{7}{8}s_k = f(w) - \frac{7}{8}s_k \leq f(c) - \frac{3}{4}s_k,$$

where the last inequality holds since $|f(c) - f(w)| \leq |c - w| \leq qt/8 \leq s_k/8$. Now consider $L = L(y) \subset \Omega \cap \mathcal{R}$ and let L' be the vertical line segment inside $B(c, qt/8M) \subset \Omega^- \cap \mathcal{R}$ through c with length $qt/4M$. By (3.14), $f(L') \subset f(L)$. Hence $\pi(L) = \pi(y)$ and $\pi(L') = \pi(c)$ belong to K and $f(L) \cap f(L') = f(L')$ is a line segment of length $\geq \lceil 4^{n-1} \beta^{-1} N \rceil^{-1} t_K$. By Lemma 3.3, the cube $K \subset \Lambda$. We have proved that if $C_1 \geq 16\alpha/q$ and there exists a cube $\mathcal{Q} \subset I$ of side length qt such that $\lambda(\mathcal{Q}) \leq \lambda(\frac{1}{2}\mathcal{Q}) + s_k/8$, then I contains a cube K in $\Lambda \cup F_k^c$ of side length $\geq \lceil 4^{n-1} \beta^{-1} N \rceil^{-1} qt/4M$.

To finish the lemma, set $m_0 = \lceil 32^{n-1}\beta^{-1}N \rceil$ and consider the descending sequence of cubes $I \supset J_1 \supset \dots \supset J_{m_0}$ with the same center as I but with side lengths $t, t/2, \dots, t/2^{m_0}$. There are three alternatives. First if $J_m \cap \pi(T) = \emptyset$ for some $1 \leq m \leq m_0$, then $J_m \cap F_k = \emptyset$ and we set $J = J_m$. Otherwise we know $\lambda(J_m)$ is defined for every m . Second suppose that for each $m < m_0$, $\lambda(J_m) \geq \lambda(J_{m+1}) + s_k/8$. Then one can find m_0 points $y_m \in T$ such that $\pi(y_m) \in J_m \subset I$ but $|y_m - y_{m'}| \geq |f(y_m) - f(y_{m'})| \geq s_k/8 \geq t/8$ for each $m \neq m'$. The m_0 surface balls $\Delta(y_m, t/16) \subset \Delta(0, r) \cap \pi^{-1}(2I)$ are disjoint; by Lemma 2.3,

$$(3.15) \quad \begin{aligned} \mathcal{H}^{n-1}(\Delta(0, r) \cap \pi^{-1}(2I)) &\geq m_0 \beta (t/16)^{n-1} \\ &\geq 32^{n-1} N (t/16)^{n-1} = N \mathcal{H}^{n-1}(2I). \end{aligned}$$

Thus, $2I \subset \Lambda$ and we can select $J = I$. Third suppose that $\lambda(J_m) \leq \lambda(J_{m+1}) + s_k/8$ for some $1 \leq m < m_0$. Put $C_1 = 2^{m_0+3}\alpha$ (which depends only on n, N, α and β) so that $C_1 \geq 16\alpha/q$ for $q = 1/2^m$. For $\mathcal{Q} = J_m$ the argument above produces $K \subset I \cap (\Lambda \cup F_k^c)$ with side length $\geq \lceil 4^{n-1}\beta^{-1}N \rceil^{-1} q/4M$. Thus, we can set $J = K$. In the worst scenario (the last case), we found a cube $J \subset I \cap (\Lambda \cup F_k^c)$ of side length at $\geq t/2^{m_0+2n+1}\beta^{-1}NM$. Therefore, since any cube contains an admissible cube at least one-quarter of its own size, we can take $C_2 = 2^{m_0+2n+3}\beta^{-1}NM$ (which depends only on n, M, N and β). \square

Next we iterate Lemma 3.4. If h is sufficiently large, then $F_k \setminus \Lambda$ is not concentrated in any cube of size s_k .

Lemma 3.5. *For all $\delta > 0$ there exists $h_1 = h_1(n, C_1, C_2, \delta)$ such that*

$$(3.16) \quad \mathcal{H}^{n-1}(I \cap F_k \setminus \Lambda) \leq \delta \mathcal{H}^{n-1}(I)$$


whenever $h \geq h_1$, $k \geq 0$ and $I \subset I_0$ is an admissible cube of side length s_k .

Proof. Let us agree that a cube $J \subset I$ is *good* if $J \subset \Lambda \cup F_k^c$; otherwise, we call J *bad*. Let P be the smallest integer power of two at least C_2 . In round one, cover I by P^{n-1} admissible cubes of side length s_k/P . If h is sufficiently large, then Lemma 3.4 applies to I and at least one cube of size s_k/P is good and at most $P^{n-1} - 1$ cubes are bad. Round two. Cover each bad cube of size s_k/P by P^{n-1} admissible cubes of size s_k/P^2 . Applying Lemma 3.4 to each parent, we conclude the number of bad admissible cubes of size s_k/P^2 is at most $(P^{n-1} - 1)^2 = P^{2(n-1)}(1 - P^{-n+1})^2$. Repeating this procedure through round R , we conclude the number of bad admissible cubes of size s_k/P^R is at most $P^{R(n-1)}(1 - P^{-n+1})^R$. Set R to be the first positive integer such that $(1 - P^{-n+1})^R < \delta$. To invoke Lemma 3.4 for R rounds total, we needed $s_k/P^q \geq C_1 h^{-1} s_k$ for each $q < R$. Thus, if $h \geq h_1 = C_1 P^{R-1}$, then the number of bad admissible cubes J of size s_k/P^R is at most $\delta P^{R(n-1)}$. It follows that

$$(3.17) \quad \begin{aligned} \mathcal{H}^{n-1}(I \cap F_k \setminus \Lambda) &\leq \sum_{J \subset I} \mathcal{H}^{n-1}(J \cap F_k \setminus \Lambda) \\ &\leq \delta P^{R(n-1)} (s_k/P^R)^{n-1} = \delta \mathcal{H}^{n-1}(I) \end{aligned}$$

as desired. \square

For each $k \geq 0$, call $(I_{k,j})_j$ the sequence of all admissible cubes of side length s_k that meet $F_k \setminus \Lambda$. In order to control the sum in (3.6), we associate a piece of surface $S(I_{k,j})$ to every cube $I_{k,j}$ and then study the size and overlap of the $S(I_{k,j})$. Here is the construction:

FIGURE 6. A piece of surface $S(I_{k,j})$

Fix a large constant $\zeta \gg 1$ to be chosen later. Let $I = I_{k,j}$ be an admissible cube of side length s_k that meets $F_k \setminus \Lambda$. Let $y = y(I)$ and $z = z(I)$ be any two points of T_E such that $\pi(y) \in I \cap (F_k \setminus \Lambda)$, such that $z \in y + \mathcal{C}$, and such that $s_k \leq f(z) - f(y) \leq s_{k-1}$. Let $c = c(I)$ be any non-tangential point $c = A^-(z, s_k/\zeta)$ of z in Ω^- . Then $|c - z| \leq s_k/\zeta$ and $B(c, s_k/\zeta M) \subset \Omega^- \cap B(z, 2s_k/\zeta)$. Furthermore, $B(c, s_k/\zeta M) \subset \mathcal{R}$ if we select $\zeta \geq 8$ (compare points in the ball to $z \in T_E$). Assign $D = D(I)$ to be the $(n-1)$ -dimensional disk with center c and radius $s_k/2\zeta M$ that is parallel to I_0 . We define $S = S(I)$ to be the set of all points w such that

- (1) $w \in \partial\Omega \cap \mathcal{R}$ or $w \in \pi(D) \times \{-b_n\}$,
- (2) $\pi(w) \in \pi(D)$,
- (3) $f(w) \leq f(c) - s_k/\zeta M$, and
- (4) the open vertical segment joining w to $\hat{w} = (\pi(w), f(c) - s_k/\zeta M)$ does not intersect $\partial\Omega$.

Including the extra $(n-1)$ cube $\pi(D) \times \{-b_n\}$ in the definition of S ensures that the projection $\pi(S) = \pi(D)$ is a disk of radius $s_k/2\zeta M$.

Lemma 3.6. *Assume that $h \geq 4\alpha$ and $\zeta \geq 8$. Then $\pi(S(I_{k,j})) \subset 2I_{k,j}$.*

Proof. Let $I = I_{k,j}$ and write y, z, c, D and S for the data associated to I . Then $\pi(y) \in I$ and $|\pi(z) - \pi(y)| \leq h^{-1}(f(z) - f(y)) \leq s_{k-1}/h = \alpha s_k/h$. Since the disk $\pi(S) = \pi(D)$ of radius $s_k/2\zeta M$ is centered at $\pi(c)$ and $|\pi(c) - \pi(z)| \leq s_k/\zeta$, we get that, for all $x \in \pi(S)$,

$$(3.18) \quad \text{dist}(x, I) \leq |x - \pi(c)| + |\pi(c) - \pi(z)| + |\pi(z) - \pi(y)| \leq \frac{s_k}{2\zeta M} + \frac{s_k}{\zeta} + \frac{\alpha s_k}{h}.$$

Hence, $\text{dist}(x, I) \leq s_k/2$ and $x \in 2I$, if we require that $\zeta \geq 8$ and $h \geq 4\alpha$. \square

Set $S_k = \bigcup_j S(I_{k,j})$. The measure of $F_k \setminus \Lambda$ is controlled by the measure of $\pi(S_k)$.

Lemma 3.7. *There exists $h_2 = h_2(n, M, \gamma, \varepsilon, C_1, C_2, \zeta)$ such that if $h \geq \max(h_2, 4\alpha)$, then for each $k \geq 0$,*

$$(3.19) \quad \mathcal{H}^{n-1}(F_k \setminus \Lambda) \leq \frac{\varepsilon}{2} \left(\gamma + \frac{1}{M\sqrt{n-1}} \right)^{-1} \mathcal{H}^{n-1}(\pi(S_k)).$$

Proof. Let $h_2 = h_1(n, C_1, C_2, \delta)$ from Lemma 3.5 where

$$(3.20) \quad \delta = \frac{\varepsilon}{2} \left(\gamma + \frac{1}{M\sqrt{n-1}} \right)^{-1} \times \frac{\omega_{n-1}}{(6\zeta M)^{n-1}}.$$

If p is the number of admissible cubes of size s_k that meet $F_k \setminus \Lambda$, then

$$(3.21) \quad \mathcal{H}^{n-1}(F_k \setminus \Lambda) \leq \sum_j \mathcal{H}^{n-1}(I_{k,j} \cap F_k \setminus \Lambda) \leq p\delta s_k^{n-1}.$$

Fix a cube $I = I_{k,j}$. By Lemma 3.6, $\pi(S(I)) = \pi(D(I))$ is a disk of radius $s_k/2\zeta M$ that is contained in $2I$. Hence $\mathcal{H}^{n-1}(\pi(S_k) \cap 2I) \geq \omega_{n-1}(s_k/2\zeta M)^{n-1}$. Because the cubes $(2I_{k,j})_j$ have bounded overlap (each $x \in \mathbb{R}^{n-1}$ lies in at most 3^{n-1} cubes),

$$(3.22) \quad p\omega_{n-1}(s_k/2\zeta M)^{n-1} \leq \sum_j \mathcal{H}^{n-1}(\pi(S_k) \cap 2I_{k,j}) \leq 3^{n-1} \mathcal{H}^{n-1}(\pi(S_k)).$$

(This step uses the fact that the sets $\pi(S_k)$ and $\pi(S_k) \cap 2I_{k,j}$ are measurable.) Thus, $\mathcal{H}^{n-1}(F_k \setminus \Lambda) \leq \delta(6\zeta M)^{n-1} \mathcal{H}^{n-1}(\pi(S_k))/\omega_{n-1}$, explaining our choice of δ . \square

By choosing good parameters, the pieces of surface S_k can be made disjoint!

Lemma 3.8. *We can find $\alpha \geq 4M\sqrt{n-1}$ depending only on n and M , and find $\zeta \geq 8$ depending only on n , N and β such that $\overline{S_k} \cap \overline{S_{k'}} = \emptyset$ for all $k \neq k'$ whenever $h \geq \alpha\zeta$.*

Proof. To start assume that $h \geq 4\alpha$ and $\zeta \geq 8$. Then the pieces of surface $(S(I_{k,j}))_j$ have finite overlap by Lemma 3.6. Hence $\overline{S_k} = \bigcup_j \overline{S(I_{k,j})}$ for each k .

Let cubes $I = I_{k,j}$ and $I' = I_{k',j'}$ be given. We shall write y, z, c, D and S for the data associated to I and write y', z', c', D' and S' for the data associated to I' . Suppose to get a contradiction that $k \geq k' + 1$ and $\overline{S} \cap \overline{S'} \neq \emptyset$. Then we can find $w \in S$ and $w' \in S'$ such that

$$(3.23) \quad |w - w'| \leq \frac{s_{k'}}{4\zeta M}.$$

Let L_w denote the vertical line segment joining w to $\hat{w} = (\pi(w), f(c) - s_k/\zeta M)$. Also write $L_y = L(y) \subset \Omega$ for the vertical line segment over y in \mathcal{T} and set $B' = B(c', s_{k'}/\zeta M) \subset \Omega^-$.

Our first claim is that $\pi(y) \in \pi(B')$ and $f(y) > f(c')$ for certain choices h, α and ζ . Indeed, since $\pi(w) \in \pi(S)$,

$$(3.24) \quad |\pi(w) - \pi(y)| \leq |\pi(w) - \pi(c)| + |\pi(c) - \pi(z)| + |\pi(z) - \pi(y)| \leq \frac{s_k}{2\zeta M} + \frac{s_k}{\zeta} + \frac{\alpha s_k}{h}.$$

If we select $h \geq \alpha\zeta$, then (3.24) implies that $|\pi(w) - \pi(y)| \leq 3s_k/\zeta$. Hence,

$$(3.25) \quad \begin{aligned} |\pi(y) - \pi(c')| &\leq |\pi(y) - \pi(w)| + |\pi(w) - \pi(w')| + |\pi(w') - \pi(c')| \\ &\leq \frac{3s_k}{\zeta} + \frac{s_{k'}}{4\zeta M} + \frac{s_{k'}}{2\zeta M} \leq \frac{3s_{k'}}{\alpha\zeta} + \frac{s_{k'}}{4\zeta M} + \frac{s_{k'}}{2\zeta M}, \end{aligned}$$

where $s_k \leq s_{k'}/\alpha$ since $k \geq k' + 1$. Thus, $|\pi(y) - \pi(c')| \leq s_{k'}/\zeta M$ and $\pi(y) \in \pi(B')$, provided that $\alpha \geq 12M$. If $f(y) \leq f(c')$, then $\pi(y) \in \pi(B')$ implies that $L_y \subset \Omega$ intersects $B' \cap f^{-1}(c') \subset \Omega^-$, which is absurd. Therefore, $f(y) > f(c')$, as claimed.

Next we claim L_w is contained in Ω^- . On one hand the upper endpoint \hat{w} of L_w satisfies

$$(3.26) \quad f(\hat{w}) \geq f(z) - \frac{2s_k}{\zeta} \geq f(z) - \frac{s_k}{4} \geq f(y) + \frac{3s_k}{4} > f(c')$$

since $\zeta \geq 8$. On the other hand the lower endpoint of L_w satisfies,

$$(3.27) \quad f(w) \leq f(w') + \frac{s_{k'}}{4\zeta M} \leq f(c') - \frac{3s_{k'}}{4\zeta M} < f(c').$$

Thus, since $\pi(w) \in \pi(B')$ (in fact $|\pi(w) - \pi(c')| \leq \frac{3}{4}s_{k'}/\zeta M$), the line segment L_w intersects $B' \cap f^{-1}(c') \subset \Omega^-$. But L_w does not intersect $\partial\Omega$ (by definition of S), so $L_w \subset \Omega^-$.

Finally, since $f(\hat{w}) \geq f(y) + 3s_k/4$ and $f(y) > f(w)$, we know $f(L_w) \cap f(L_y)$ is an interval of length at least $3s_k/4$. We previously showed that $|\pi(w) - \pi(y)| \leq 3s_k/\zeta$ if $h \geq \zeta\alpha$. Thus, L_w and L_y lie over a cube J of side length $\lceil 4^{n-1}\beta^{-1}N \rceil^{-1}3s_k/4$ if $\zeta \geq 4\lceil 4^{n-1}\beta^{-1}N \rceil$. By Lemma 3.3, $\pi(y) = \pi(L_y) \in J \subset \Lambda$. This is a contradiction, because $\pi(y) \notin \Lambda$ by the definition of y . An examination of the conditions on parameters assumed above reveals the lemma holds with $\zeta = 4\lceil 4^{n-1}\beta^{-1}N \rceil$, $\alpha = \max(4M\sqrt{n-1}, 12M)$ and $h \geq \alpha\zeta$. \square

We are ready to conclude. Use Lemma 3.8 to pick α and ζ , and set $h = \max(h_0, h_2, \alpha\zeta)$. Then the pieces of surface $\overline{S_k} \subset \mathcal{S} := \Delta(Q, r) \cup (2I_0 \times \{-b_n\})$ are disjoint and measurable. Thus

$$(3.28) \quad \sum_{k=0}^{\infty} \mathcal{H}^{n-1}(\pi(S_k)) \leq \sum_{k=0}^{\infty} \mathcal{H}^{n-1}(\overline{S_k}) \leq \mathcal{H}^{n-1}(\mathcal{S}) \leq \left(\gamma + \frac{1}{M\sqrt{n-1}} \right) r^{n-1}.$$

Using Lemma 3.7, we conclude

$$(3.29) \quad \sum_{k=0}^{\infty} \mathcal{H}^{n-1}(F_k \setminus \Lambda) \leq \frac{\varepsilon}{2} \left(\gamma + \frac{1}{M\sqrt{n-1}} \right)^{-1} \sum_{k=0}^{\infty} \mathcal{H}^{n-1}(\pi(S_k)) \leq \frac{\varepsilon}{2} r^{n-1}.$$

Therefore, (3.6) holds and Proposition 2.8 is established.

4. HARMONIC MEASURE ON NTA DOMAINS

In this section we prove Theorem 1.2 on the absolute continuity of harmonic measure. An NTA domain is a corkscrew domain (studied in §2 above) that also admits a Harnack chain condition. The class of NTA domains was introduced by Jerison and Kenig [8]. Given $X_1, X_2 \in \Omega$ a *Harnack chain* from X_1 to X_2 is a sequence of open balls in Ω such that the first ball contains X_1 , the last ball contains X_2 , and consecutive balls intersect.

Definition 4.1. A domain $\Omega \subset \mathbb{R}^n$ satisfies the *Harnack chain condition* with constants $M > 1$ and $R > 0$ if for every $Q \in \partial\Omega$ and $0 < r < R$ when $X_1, X_2 \in \Omega \cap B(Q, r)$ satisfy

$$(4.1) \quad \min_{j=1,2} \text{dist}(X_j, \partial\Omega) > \varepsilon \quad \text{and} \quad |X_1 - X_2| < 2^k \varepsilon$$

then there is a Harnack chain from X_1 to X_2 of length Mk such that the diameter of each ball is bounded below by $M^{-1} \min_{j=1,2} \text{dist}(X_j, \partial\Omega)$.

Definition 4.2. A domain $\Omega \subset \mathbb{R}^n$ is *non-tangentially accessible* or *NTA* if there exist $M > 1$ and $R > 0$ such that (i) Ω satisfies the corkscrew and Harnack chain conditions, (ii) $\mathbb{R}^n \setminus \Omega$ satisfies the corkscrew condition. If $\partial\Omega$ is unbounded, then we require $R = \infty$.

The exterior corkscrew condition guarantees that an NTA domain $\Omega \subset \mathbb{R}^n$ is regular for the Dirichlet problem; i.e. for every $f \in C_c(\partial\Omega)$ there exists $u \in C^2(\Omega) \cap C(\overline{\Omega})$ such that $\Delta u = 0$ in Ω and $u = f$ on $\partial\Omega$. Thus the maximum principle and Riesz representation theorem yield a family of Borel regular probability measures $\{\omega^X\}_{X \in \Omega}$ on $\partial\Omega$ such that

$$(4.2) \quad u(X) = \int_{\partial\Omega} f(Q) d\omega^X(Q)$$

is the unique harmonic extension of $f \in C_c(\partial\Omega)$. We call ω^X the *harmonic measure* of Ω with pole at X . Because $\omega^{X_1} \ll \omega^{X_2} \ll \omega^{X_1}$ for any $X_1, X_2 \in \Omega$ (by Harnack's inequality), it makes sense to discuss null sets of harmonic measure $\omega = \omega^{X_0}$ with respect to some fixed pole $X_0 \in \Omega$ far away from the boundary.

The special feature of harmonic measure on NTA domains (versus corkscrew domains) that we need below is the the following localization property.

Lemma 4.3 ([8] (4.18)). *Let $\Omega \subset \mathbb{R}^n$ be NTA. There exists $C = C(n, M) > 0$ with the following property. Assume $X_0 \in \Omega \setminus B(Q, 2r)$ for some $Q \in \partial\Omega$ and $r < R/2$. Then for every non-tangential point $a = A^+(Q, r)$ and every Borel set $E \subset \Delta(Q, r)$,*

$$(4.3) \quad C^{-1}\omega^a(E) \leq \frac{\omega(E)}{\omega(\Delta(Q, r))} \leq C\omega^a(E).$$

Remark 4.4. In Definition 4.2 we allow our NTA domains $\Omega \subset \mathbb{R}^n$ to be either bounded or unbounded. The proof of Lemma 4.3 for bounded domains in [8] carries through to the unbounded case without modification.

At every boundary point of finite lower density there is a shrinking sequence of scales on which the harmonic measure and the surface measure are comparable in the sense of (1.2). The proof of Proposition 4.5 below follows the same structure of David and Jerison's proof of Theorem 2 in [4]; however, we keep careful track of the constants appearing from the Lipschitz approximations of the domain (Theorem 2.4 and Lemma 2.13). The required technical tools are the localization principle for harmonic measure (Lemma 4.3) and the maximum principle for harmonic functions.

Proposition 4.5. *There exist constants $0 < \delta < 1$ and $0 < \varepsilon < 1$ depending only on n, M and γ with the following property. Let $\Omega \subset \mathbb{R}^n$ be an NTA domain with constants $M > 1$ and $R > 0$. If $\liminf_{r \downarrow 0} \mathcal{H}^{n-1}(\Delta(Q, r))/r^{n-1} < \gamma < \infty$, then there is a sequence of positive numbers $r_i < R$ such that $\lim_{i \rightarrow \infty} r_i = 0$ and for every Borel set $E \subset \Delta(Q, r_i)$:*

$$(4.4) \quad \omega(E) \leq \delta \omega(\Delta(Q, r_i)) \Rightarrow \mathcal{H}^{n-1}(E) \leq \varepsilon \mathcal{H}^{n-1}(\Delta(Q, r_i)),$$

$$(4.5) \quad \mathcal{H}^{n-1}(E) \leq \delta \mathcal{H}^{n-1}(\Delta(Q, r_i)) \Rightarrow \omega(E) \leq \varepsilon \omega(\Delta(Q, r_i)).$$

Proof. Let $\Omega \subset \mathbb{R}^n$ be NTA with constants $M > 1$ and $R > 0$ and assume that $Q \in \partial\Omega$ satisfies $\liminf_{r \downarrow 0} \mathcal{H}^{n-1}(\Delta(Q, r))/r^{n-1} < \gamma < \infty$. Then there exists a sequence of positive numbers $r_i < R$ decreasing to zero such that $\mathcal{H}^{n-1}(\Delta(Q, r_i)) \leq \gamma r_i^{n-1}$. Let $\psi = \psi(n, M)$ and $\eta = \eta(n, M, \gamma)$ be the constants given by Theorem 2.4 and Lemma 2.13. By passing

to a subsequence of r_i if necessary, we may assume that the pole of ω lies outside $B(Q, 2r_i)$ for all i (so that we can invoke the localization principle for NTA domains).

Fix r_i and pick any non-tangential point $a = A^+(Q, r_i/2)$ of Ω . Let ω^a denote harmonic measure of Ω with pole at a and let $\Delta = \Delta(Q, r_i)$. By the localization principle,

$$(4.6) \quad C^{-1}\omega^a(E) \leq \frac{\omega(E)}{\omega(\Delta)} \leq C\omega^a(E) \quad \text{for every Borel set } E \subset \Delta,$$

where the constant $C > 1$ only depends on the NTA constants of Ω . By Theorem 2.4 there is a Lipschitz domain $\Omega_L \subset \mathbb{R}^n$ such that $a \in \Omega_L \subset \Omega \cap B(Q, r_i)$ and $F = \partial\Omega_L \cap \partial\Omega$ satisfies $\mathcal{H}^{n-1}(F) \geq \psi r_i^{n-1}$. Let ω_L^a denote the harmonic measure of Ω_L with pole at a .

Assume that $E \subset \Delta$ is Borel and $\omega(E) < \delta\omega(\Delta)$. By the maximum principle and (4.6),

$$(4.7) \quad \omega_L^a(E \cap F) \leq \omega^a(E \cap F) \leq \omega^a(E) \leq C\delta.$$

If $C\delta \leq \eta$, then (2.26) and (4.7) imply that $\mathcal{H}^{n-1}(E \cap F) \leq (\psi/2)r_i^{n-1}$. Hence

$$(4.8) \quad \mathcal{H}^{n-1}(F \setminus E) = \mathcal{H}^{n-1}(F) - \mathcal{H}^{n-1}(E \cap F) \geq \psi r_i^{n-1} - (\psi/2)r_i^{n-1} = (\psi/2)r_i^{n-1}.$$

It follows that

$$(4.9) \quad \begin{aligned} \mathcal{H}^{n-1}(E) &= \mathcal{H}^{n-1}(\Delta) - \mathcal{H}^{n-1}(\Delta \setminus E) \leq \mathcal{H}^{n-1}(\Delta) - \mathcal{H}^{n-1}(F \setminus E) \\ &\leq \mathcal{H}^{n-1}(\Delta) - \frac{\psi}{2}r_i^{n-1} = \mathcal{H}^{n-1}(\Delta) - \frac{\psi}{2\gamma}\gamma r_i^{n-1} \\ &\leq \mathcal{H}^{n-1}(\Delta) - \frac{\psi}{2\gamma}\mathcal{H}^{n-1}(\Delta) = \left(1 - \frac{\psi}{2\gamma}\right)\mathcal{H}^{n-1}(\Delta). \end{aligned}$$

Thus (4.4) holds for all $0 < \delta \leq \eta/C$ and for all $1 - \psi/2\gamma \leq \varepsilon < 1$.

Now assume $E \subset \Delta$ satisfies $\mathcal{H}^{n-1}(E) \leq \delta\mathcal{H}^{n-1}(\Delta) \leq \delta\gamma r_i^{n-1}$. If $\delta\gamma \leq \psi/4$, then $\mathcal{H}^{n-1}(F \setminus E) = \mathcal{H}^{n-1}(F) - \mathcal{H}^{n-1}(E \cap F) \geq (3\psi/4)r_i^{n-1}$. Hence the contrapositive of (2.26) implies that $\omega_L^a(F \setminus E) > \eta$. By the maximum principle and (4.6), $\omega(F \setminus E) \geq (\eta/C)\omega(\Delta)$. We conclude

$$(4.10) \quad \omega(E) = \omega(\Delta) - \omega(\Delta \setminus E) \leq \omega(\Delta) - \omega(F \setminus E) \leq (1 - \eta/C)\omega(\Delta).$$

Thus (4.5) holds for all $0 < \delta \leq \psi/4\gamma$ and for all $1 - \eta/C \leq \varepsilon < 1$. Therefore the constants

$$(4.11) \quad \delta = \min\{\eta/C, \psi/4\gamma\} \quad \text{and} \quad \varepsilon = 1 - \delta$$

which depend only on n , M and κ suffice. \square

To stitch together estimates in Proposition 4.5 at different points, we use a Vitali type covering lemma for Radon measures in \mathbb{R}^n .

Theorem 4.6 ([13] Theorem 2.8). *Let μ be a Radon measure on \mathbb{R}^n , $A \subset \mathbb{R}^n$ and \mathcal{B} a family of closed balls such that each point of A is the center of arbitrarily small balls; i.e.,*

$$(4.12) \quad \inf\{r : B(x, r) \in \mathcal{B}\} = 0 \quad \text{for all } x \in A.$$

Then there are disjoint balls $B_i \in \mathcal{B}$ such that

$$(4.13) \quad \mu\left(A \setminus \bigcup_i B_i\right) = 0.$$

We now establish the main theorem. Recall: *Let $\Omega \subset \mathbb{R}^n$ be NTA. Then the set*

$$(4.14) \quad A = \left\{ Q \in \partial\Omega : \liminf_{r \downarrow 0} \frac{\mathcal{H}^{n-1}(\Delta(Q, r))}{r^{n-1}} < \infty \right\}$$

is $(n-1)$ -rectifiable and $\omega \ll A \ll \sigma \ll A \ll \omega \ll A$.

Proof of Theorem 1.2. The set A is $(n-1)$ -rectifiable by Corollary 2.6. Define the set

$$(4.15) \quad A_k = \left\{ Q \in \partial\Omega : \liminf_{r \downarrow 0} \frac{\mathcal{H}^{n-1}(\Delta(Q, r))}{r^{n-1}} < k \right\}$$

for each integer $k \geq 1$. Then $A = \bigcup_{k=1}^{\infty} A_k$ and to show $\omega \ll A \ll \sigma \ll A \ll \omega \ll A$ we may prove that $\omega(E) = 0$ if and only if $\mathcal{H}^{n-1}(E) = 0$ for every $k \geq 1$ and compact set $E \subset A_k$. (It is enough to take E compact, because $\mathcal{H}^{n-1} \ll A_k$ is Radon.)

Let $E \subset A_k$ be an arbitrary compact set and assume that $\omega(E) = 0$. Since

$$(4.16) \quad \liminf_{r \downarrow 0} \mathcal{H}^{n-1}(\Delta(Q, r))/r^{n-1} < k$$

for all $Q \in E$, Proposition 4.5 implies there exists constants $0 < \delta < 1$ and $0 < \varepsilon < 1$ such that (4.4) and (4.5) hold for each $Q \in E$ (along some sequence $r_i \downarrow 0$ depending on Q). Let U be any (relatively) open set $U \subset \partial\Omega$ such that $E \subset U$. The family $\mathcal{F} = \{\Delta(Q, r_i)\}$ of all closed balls with center $Q \in E$ and radii r_i satisfying $\Delta(Q, r_i) \subset U$ and (4.4) is a fine cover of E (in the sense of (4.12)). By Theorem 4.6, there exists a disjoint sequence Δ_i of disks in \mathcal{F} such that

$$(4.17) \quad \mathcal{H}^{n-1} \left(E \setminus \bigcup_i \Delta_i \right) = 0.$$

Since $\omega(E \cap \Delta_i) = 0 \leq \delta \omega(\Delta_i)$ for each i , $\mathcal{H}^{n-1}(E \cap \Delta_i) \leq \varepsilon \mathcal{H}^{n-1}(\Delta_i)$ for each i by (4.4). Thus, by (4.17),

$$(4.18) \quad \mathcal{H}^{n-1}(E) = \sum_i \mathcal{H}^{n-1}(E \cap \Delta_i) \leq \varepsilon \sum_i \mathcal{H}^{n-1}(\Delta_i) \leq \varepsilon \mathcal{H}^{n-1}(U).$$

Because $U \supset E$ was an arbitrary open set, by the outer regularity of Radon measures, $\mathcal{H}^{n-1}(E) \leq \varepsilon \mathcal{H}^{n-1}(E)$. But $\mathcal{H}^{n-1}(E) < \infty$ (since E is compact) and $0 < \varepsilon < 1$. Therefore, $\mathcal{H}^{n-1}(E) = 0$ whenever $\omega(E) = 0$.

If $E \subset A_k$ is a compact set such that $\mathcal{H}^{n-1}(E) = 0$, then the same argument with the roles of ω and \mathcal{H}^{n-1} reversed and (4.5) in place of (4.4) shows $\omega(E) = 0$. This completes the proof of absolute continuity on A . \square

5. HAUSDORFF DIMENSION AND WOLFF SNOWFLAKES

We now present two applications of Theorem 1.2 to the dimension of harmonic measure. Let $\dim E$ denote Hausdorff dimension of a set $E \subset \mathbb{R}^n$. The (*upper*) *Hausdorff dimension* of harmonic measure,

$$(5.1) \quad \mathcal{H}\text{-dim } \omega = \inf \{ \dim E : E \subset \partial\Omega \text{ is Borel and } \omega(E) = 1 \},$$

is the smallest dimension of a set with full harmonic measure. In [12] Makarov determined $\mathcal{H}\text{-dim } \omega = 1$ for every simply connected planar domain (independent of the Hausdorff dimension of the boundary); moreover, $\omega \ll \mathcal{H}^s$ for all $s < 1$ and $\omega \perp \mathcal{H}^t$ for all $t > 1$. Higher dimensions display different behavior. Wolff [16] constructed NTA domains $\Omega \subset \mathbb{R}^3$

such that $\mathcal{H}\text{-dim } \omega > 2$ and other NTA domains $\Omega \subset \mathbb{R}^3$ such that $\mathcal{H}\text{-dim } \omega < 2$. Extending this construction, Lewis, Verchota and Vogel [11] built 2-sided NTA domains $\Omega \subset \mathbb{R}^n$ (i.e. $\Omega^+ = \Omega$ and $\Omega^- = \mathbb{R}^n \setminus \overline{\Omega}$ are both NTA) called *Wolff snowflakes* such that

- (1) $\mathcal{H}\text{-dim } \omega^+ > n - 1$ and $\mathcal{H}\text{-dim } \omega^- > n - 1$,
- (2) $\mathcal{H}\text{-dim } \omega^+ > n - 1$ and $\mathcal{H}\text{-dim } \omega^- < n - 1$,
- (3) $\mathcal{H}\text{-dim } \omega^+ < n - 1$ and $\mathcal{H}\text{-dim } \omega^- < n - 1$.

Here ω^+ denotes harmonic measure on the interior Ω^+ and ω^- denotes harmonic measure on the exterior Ω^- of Ω . While surface measure $\sigma = \mathcal{H}^{n-1} \llcorner \partial\Omega$ is clearly infinite for Wolff snowflakes of type (1) or (2), the same is not apparent for snowflakes of type (3). This is the first application of Theorem 1.2: every Wolff snowflake has infinite surface measure. In fact, if the dimension of harmonic measure on a (1-sided) NTA domain is small, then the surface measure is infinite at all locations and scales.

Theorem 5.1. *Let $\Omega \subset \mathbb{R}^n$ be NTA. If $\mathcal{H}\text{-dim } \omega < n - 1$, then σ is locally infinite; i.e., $\mathcal{H}^{n-1}(\Delta(Q, r)) = \infty$ for every $Q \in \partial\Omega$ and $r > 0$.*

Proof. Assume that $\Omega \subset \mathbb{R}^n$ is NTA and $\mathcal{H}\text{-dim } \omega < n - 1$. Then there exists a Borel set $E \subset \partial\Omega$ such that $\dim E < n - 1$ and $\omega(\partial\Omega \setminus E) = 0$. Suppose for contradiction that $\mathcal{H}^{n-1}(\Delta(Q, r)) < \infty$ for some $Q \in \partial\Omega$ and $r > 0$. Then harmonic measure and surface measure have the same null sets on $\Delta(Q, r) \cap A$ by Theorem 1.2. On one hand,

$$(5.2) \quad \omega(\Delta(Q, r) \cap A) = \omega(E \cap \Delta(Q, r) \cap A) = 0$$

since $\mathcal{H}^{n-1}(E) = 0$. On the other hand,

$$(5.3) \quad \omega(\Delta(Q, r) \cap A) > 0$$

since $\mathcal{H}^{n-1}(\Delta(Q, r) \cap A) = \mathcal{H}^{n-1}(\Delta(Q, r)) > 0$ by Lemma 2.3. The fallacy is apparent. We conclude that $\mathcal{H}^{n-1}(\Delta(Q, r)) = \infty$ for every $Q \in \partial\Omega$ and $r > 0$. \square

In [9] Kenig, Preiss and Toro used tangent measures of harmonic measure to compute $\mathcal{H}\text{-dim } \omega^+ = \mathcal{H}\text{-dim } \omega^- = n - 1$ on every 2-sided NTA domain $\Omega \subset \mathbb{R}^n$ ($n \geq 3$) with $\omega^+ \ll \omega^- \ll \omega^+$. (Thus the interior and exterior harmonic measures on Wolff snowflakes are mutually singular.) Using Theorem 1.1, we obtain a similar result for (1-sided) NTA domains of locally finite perimeter.

Theorem 5.2. *Let $\Omega \subset \mathbb{R}^n$ be NTA. If $\mathcal{H}^{n-1} \llcorner \partial\Omega$ is Radon, then $\mathcal{H}\text{-dim } \omega = n - 1$.*

Proof. If $E \subset \partial\Omega$ has dimension $t < n - 1$ then $\mathcal{H}^{n-1}(\partial\Omega \setminus E) > 0$. Hence, $\omega(\partial\Omega \setminus E) > 0$, because $\sigma \ll \omega$ by Theorem 1.1. Since no set $E \subset \partial\Omega$ of dimension $t < n - 1$ has full harmonic measure, we get $\mathcal{H}\text{-dim } \omega \geq n - 1$. Conversely, $\mathcal{H}\text{-dim } \omega \leq \dim \partial\Omega = n - 1$, since $\mathcal{H}^{n-1} \llcorner \partial\Omega$ is Radon. \square

REFERENCES

- [1] B. Bennewitz and J. Lewis, *On weak reverse Hölder inequalities for nondoubling harmonic measures*, Complex Var. Theory Appl. **49** (2004), no. 7–9, 571–582.
- [2] R. R. Coifman and C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, Studia Math. **51** (1974), 241–250.
- [3] B. Dahlberg, *Estimates of harmonic measure*, Arch. Rational Mech. Anal. **65** (1977), no. 3, 275–288.
- [4] G. David and D. Jerison, *Lipschitz approximation to hypersurfaces, harmonic measure, and singular integrals*, Indiana Univ. Math. J. **39** (1990), no. 3, 831–845.

- [5] G. David and S. Semmes, *Analysis of and on uniformly rectifiable sets*, Mathematical Surveys and Monographs, vol. 38, American Mathematical Society, Providence, R.I., 1993.
- [6] J. Garnett and D. Marshall, *Harmonic measure*, New Mathematical Monographs, vol. 2, Cambridge University Press, Cambridge, 2005.
- [7] D. Jerison and C. Kenig, *An identity with applications to harmonic measure*, Bull. Amer. Math. Soc. (N.S.) **2** (1980), no. 3, 447–451.
- [8] ———, *Boundary behavior of harmonic functions in non-tangentially accessible domains*, Adv. in Math. **46** (1982), no. 1, 80–147.
- [9] C. Kenig, D. Preiss, and T. Toro, *Boundary structure and size in terms of interior and exterior harmonic measures in higher dimension*, J. Amer. Math. Soc. **22** (2009), no. 3, 771–796.
- [10] M. Lavrentiev, *Boundary problems in the theory of univalent functions*, Math. Sb. (N.S.) **1** (1936), 815–845; English transl., Amer. Math. Soc. Transl. (2) **32** (1963), 1–35.
- [11] J. Lewis, G. C. Verchota, and A. Vogel, *On Wolff snowflakes*, Pacific J. Math. **218** (2005), no. 1, 139–166.
- [12] N. G. Makarov, *On the distortion of boundary sets under conformal mappings*, Proc. London Math. Soc. (3) **51** (1985), no. 2, 369–384.
- [13] P. Mattila, *Geometry of sets and measures in Euclidean spaces: Fractals and rectifiability*, Cambridge Series in Advanced Mathematics, vol. 44, Cambridge University Press, Cambridge, 1995.
- [14] F. Riesz and M. Riesz, *Über Radwerte einer analytischen Functionen*, Compte rendu du quatrième Congrès des Mathématiciens Scandinaves: tenu à Stockholm du 30 août au 2 Septembre 1916, Malmö, 1955, pp. 27–44.
- [15] S. Semmes, *Analysis vs. geometry on a class of rectifiable hypersurfaces in \mathbb{R}^n* , Indiana Univ. Math. J. **39** (1990), no. 4, 1005–1035.
- [16] T. Wolff, *Counterexamples with harmonic gradients in \mathbb{R}^3* , Essays on Fourier analysis in honor of Elias M. Stein, Princeton Mathematical Series, vol. 42, Princeton University Press, Princeton, NJ, 1995, pp. 321–384.
- [17] W. Ziemer, *Some remarks on harmonic measure in space*, Pacific J. Math. **55** (1974), 629–637.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195-4530
E-mail address: mbadger@math.washington.edu